



F I G. 2 (A)




F I G. 2 (B)




F I G. 3 (A)



F I G. 3. (B)



F I G. 3 (C)



000000-4400

FIG. 4 (A)




FIG. 4 (B)




Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The *Agrobacterium* strains were grown in the YEA medium for 24 h at 28°C. The cell concentration of the strains was adjusted to 10⁸ cells/ml. The cell suspension was mixed with the plant tissue and the transformation efficiency was determined. The results are shown as the mean ± SD of three independent experiments. The asterisk indicates a significant difference ($P < 0.05$) between the strains.






Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The *Agrobacterium* strains were grown in the YEA medium for 24 h at 28°C. The cell concentration of the strains was adjusted to 10⁸ cells/ml. The cell suspension was mixed with the plant tissue and the transformation efficiency was determined. The results are shown as the mean ± SD of three independent experiments. The asterisk indicates a significant difference ($P < 0.05$) between the strains.




F I G. 6 (A)



F I G. 6 (B)



F I G. 6 (C)



F I G. 6 (D)



[illegible]